


(ISM-2018)



615.47:616-07

61000, , « » ,
 . (057) 707-60-44, E-mail: stas69@ukr.net

This paper presents the first mathematical modeling of an oral glucose tolerance test with a double load, used for qualitative diagnosis of latent Diabetes mellitus type 2. Based on the clinical and physiological data of the carbohydrate exchange regulation system, its structural-functional minimal model in the form of the first-order differential equation with delayed argument relative to the glycemic level, that adequately reproduces the data of this test, is constructed.

2-
 (2) [1].
 (),

2
90

(2) [2].

2

2

[3].

[4],

2.

2.

$g'(t)$

$i'(t)$:

$$g'(t)_{\text{инзав}} = -\lambda i'(t), \quad (1)$$

$i'(t)$

:

$f(t)$,

$Q(t)$

$f(t)$

$g(t)$

g_b

$$y(t) = g(t) - g_b$$

:

$$i'(t)_{\text{сепр}} = \eta f(t) + \mu Q(t) + \chi y(t - \tau). \quad (3)$$

1 -

(1) (2),

$$\begin{aligned} \frac{dy}{dt} &= (1 - \alpha)f(t) - \beta y(t - 1) - \gamma y(t - \tau) - \zeta Q(t), & t \geq 0, \\ y(t) &= \phi(t), & -\tau \leq t \leq 0, \end{aligned} \quad (4)$$

t -


(

);

[5].

(4)

1 ,

$$2 (\quad .1).$$


1 -

2: - , b -

2.

2

2

2

1. : / . . , . . , . . ; . . . , . . . - . : , 2010. — 437 .
2. / - 8- . - . : - , 2016.-736 .
3. Bolie V.W. Coefficients of normal blood glucose regulation// J. Appl. Physiol. -1961. - Vol.16. - P. 783-788.
4. - : / . . , . . , . . . - : . . , 2009. - 332 .
5. . „ . „ . . . - . : - , 2015. — 880 .